

Spinal cord compression

What to look out for



Clinical Sciences Building
Acute Oncology
0161 206 2312

*Saving lives,
Improving lives*

If you have a type of cancer that has spread to a bone, you need to know about spinal cord compression and the warning signs.

If you can recognise these early warning signs and get treated quickly, permanent damage may be avoided. The aim of this booklet explains what to look out for, and what you should do if you have any warning signs.

What is spinal cord compression?

The spinal cord is a bundle of nerves that runs from the brain down the back.


The bones of the spine surround it.

The spinal cord provides nerves to the body including the bladder, arms and legs.

If you have a tumour in the bones of the spine (vertebrae) or in the tissues around the spinal cord, it can cause pressure (compression) on the spinal cord.

This can cause pain, altered sensation and weakness.

Spinal cord compression is not common and the risk of cord compression is small even if you have a secondary cancer in the spine.



What are the warning signs?

This depends on which part of the spine is affected. Pain may be present for some time before other symptoms develop. The warning signs can be any of the following:

- Pain in one area of your spine that is severe, distressing or different from your usual pain
 - Pain down your leg or arm
 - Tingling - like electric shocks - down your spine into your legs, or arms and legs
 - Severe pain in your spine which changes when you lie down or stand up, lift or strain, wakes you at night, or stops you getting to sleep
 - Pain which starts in the spine and goes around the chest or abdomen
 - New pain which gets worse and does not respond to painkillers
- Stiffness and heaviness in your legs affecting your balance and walking
 - Numbness or changed sensation in your legs - this can also affect your lower body, chest or arms
 - Weakness in your legs, or arms and legs
 - Bladder / bowel problems - loss of control (incontinence) or not being able to empty your bladder

What should I do if I develop any warning signs?

Get advice immediately: contact the hospital team where you usually go for your cancer follow-up clinics, your GP or Macmillan nurse. Describe your symptoms and explain that you are worried that you may have spinal cord compression and that you need to be seen urgently.

**Don't delay
Ring someone today even if it is at a weekend or a holiday period.**

What happens next?

- A doctor needs to examine you. If the doctor suspects that you may have spinal cord compression he will advise you to stay flat in bed and a scan of your spine will be arranged urgently. This is usually an MRI scan, but you may have a different scan if this is not suitable for you

- The doctor will prescribe steroids (dexamethasone) for you before or after the scan. Steroids reduce pressure and swelling around nerve tissue. If you are diabetic, or have had problems with steroids in the past, you should tell your doctor
- Radiotherapy, surgery or chemotherapy are all options for treatment. Your oncology team will discuss the best treatment for you with the specialist surgeons. This treatment option depends on the type of cancer, which part of the spine is affected and your general health. In any case, treatment should start as soon as possible after the scan

- If you have cord compression you will probably be admitted to your local hospital. You may need to be transferred to The Christie (Withington, Manchester or Christie@Salford) for radiotherapy or Salford Royal if you need surgery. You will have to stay flat in bed until the doctors are satisfied that your spine is stable and it is safe to start walking again. You will get help with this from the physiotherapist. If your neck or top of the (cervical) spine is affected, you will also need to wear a collar to prevent damage to the spinal cord

Spinal cord compression, if untreated or if there is a delay in treatment, can lead to permanent damage and even paralysis. Early diagnosis and treatment gives the best chance of preventing paralysis.

Remember to get help if you develop any warning signs. Seek immediate help if you have unexplained tingling, numbness or weakness in your arms and legs.

So remember:

If you have any of the above symptoms:

- Speak with a health professional, for example, your GP, Oncologist, Specialist Nurse or a Physiotherapist as soon as is practical (preferably within 24 to 48 hours)
- Tell them that you are worried about your spine and would like to be examined as soon as possible
- Bring this booklet with you
- Try to avoid bending your back

© G20020302W. Design Services
Salford Royal NHS Foundation Trust
All Rights Reserved 2020
This document **MUST NOT** be photocopied
Information Leaflet Control Policy:
Unique Identifier: NOE06(20)
Review Date: May 2022

Salford Royal **NHS**
NHS Foundation Trust

University Teaching Trust

safe • clean • personal

For further information on this leaflet, it's references and sources used, please contact **0161 206 2312**.

Copies of this leaflet are available in other formats (for example, large print or easyread) upon request. Alternative formats may also be available via www.srft.nhs.uk/for-patients/patient-leaflets/

In accordance with the Equality Act we will make reasonable adjustments to enable individuals with disabilities to access this service. Please contact the service/clinic you are attending by phone or email prior to your appointment to discuss your requirements.

Interpreters are available for both verbal and non-verbal (e.g. British Sign Language) languages, on request. Please contact the service/clinic you are visiting to request this.

If you need this information leaflet translated, please telephone:

Polish

Jeżeli potrzebne jest Państwu to tłumaczenie, proszę zadzwonić pod numer.

Urdu

اگر آپ کو اس ترجمانی کی ضرورت ہے تو مہربانی کر کے فون کریں۔

Arabic

إذا كنتم بحاجة الى تفسير او ترجمة هذا الرجاء الاتصال

Chinese

如果需要翻译，请拨打电话

Farsi

اگر به ترجمه این نیاز دارید ، لطفاً تلفن کنید

0161 206 0224

or Email:

InterpretationandTrans@srft.nhs.uk

Salford Royal operates a smoke-free policy.

For advice on stopping smoking contact the Hospital Specialist Stop Smoking Service on 0161 206 1779

This is a teaching hospital which plays an important role in the training of future doctors. The part patients play in this process is essential to make sure that we produce the right quality doctors for all of our futures. If at any time you would prefer not to have students present please inform the doctor or nurses and this will not affect your care in anyway.

Salford Royal NHS Foundation Trust
Stott Lane, Salford,
Manchester, M6 8HD
Telephone 0161 789 7373
www.srft.nhs.uk

If you would like to become a Foundation Trust Member please visit:
www.srft.nhs.uk/for-members

If you have any suggestions as to how this document could be improved in the future then please visit:
<http://www.srft.nhs.uk/for-patients>

Salford Royal is committed to producing high quality, evidence based, functional consumer health and care information. To support this we apply the principles and quality statements of The Information Standard fully to ensure that our process for producing information follows best practice.